

| <b>OUTCOME MEASURES</b>  |
|--|
| <b><i>PEDIATRIC USE ONLY:</i></b>  |
| COSI-C—Client-oriented Scale of Improvement-Children   |
| Child Outcomes Summary Form (COSF)   |
| CHILD—Children’s Home Inventory of Listening Difficulties  |
| ELF—Early Listening Function   |
| IT-MAIS—Infant –Toddler Meaningful Auditory Integration Scale  |
| Littlears Auditory Questionnaire   |
| Functional Auditory Performance Inventory  |
| MAIS—Meaningful Auditory Integration Scale   |
| Pediatric Evaluation of Disability Inventory (PEDI)  |
| Pediatric Evaluation of Disability Inventory – Computer Adaptive Test (PEDI-CAT)   |
| Pediatric Evaluation of Disability Inventory – Computer Adaptive Test (PEDI-MCAT™)   |
| Pediatric Quality of Life Inventory (PEDS-QL)  |
| Pre-Kindergarten National Outcomes Measurement System (NOMS) Functional Communication Measures <b><i>(copyrighted for use only by clinicians submitting NOMS data to ASHA and registered to participate in NOMS through their school or healthcare facility)</i></b> |
| Preschool SIFTER—Screening Instrument for Targeting Educational Risk   |
| K–12 National Outcomes Measurement System (NOMS)   |
| Early Childhood Outcomes (ECO; includes state-based variations)  |
| Wee-FIM  |
| LIFE—Listening Inventory For Education   |
| PHDANP—Pediatric Hearing Demand, Ability and Need Profile  |
| DIAL—Developmental Index of Audition and Listening   |
| FEW—Family Expectation Worksheet   |
|  |
| <b><i>ADULT USE ONLY:</i></b>  |
| Activity Measure for Post-Acute Care (AM-PAC™) [computer-based version and outpatient short-form version]  |
| ASHA Functional Assessment of Communication for Adults (ASHA FACS)   |
| American Speech-Language-Hearing Association Quality of Communication Life Scale (ASHA QCL, Paul et al., 2005)   |
| Burden of Stroke Scale (BOSS, Doyle et al., 2004)  |
| Communication Activities of Daily Living-2 (CADL-2)  |
| Communication Confidence Rating Scale for Aphasia (CCRSA, Babbitt, Heinemann, Semik, & Cherney, 2011)  |
| Communication Disability Profile (CDP, Swinburn & Byng, 2006)  |
| Communication Outcome After Stroke Scale (COAST, Long, Hesketh, Paszek, Booth, & Bowen, 2008)  |
| Communication Outcome Measures of Functional Independence: The COMFI Scale   |

|  |
|--|
| Consensus Auditory-Perceptual Evaluation of Voice (CAPE-V)   |
| Dysarthria Impact Profile  |
| Eating Assessment Tool (EAT-10)  |
| Functional Assessment Measure (FAM)  |
| Functional Independence Measure (FIM)  |
| Functional Oral Intake Scale (FOIS)  |
| Late-Life FDI™   |
| The MOS 36-item Short-Form Health Survey (SF-36)   |
| National Outcomes Measurement System (NOMS Functional Communication Measures – FCM) <i>(copyrighted for use only by clinicians submitting NOMS data to ASHA and registered to participate in NOMS through their school or healthcare facility)</i> |
| Rehabilitation Institute of Chicago Functional Assessment Scale (RIC-FAS)  |
| Reintegration to Normal Living (RNL)   |
| Stroke and Aphasia Quality of Life Scale-39 (SAQOL-39, Hilari, Byng, Lamping, & Smith, 2003)   |
| Stroke Impact Scale  |
| SWAL-QOL   |
| Therapy Outcome Measure (TOM)  |
| Voice Handicap Index (VHI)   |
|  |
| <b><i>PEDIATRIC AND ADULT:</i></b>   |
| Assessment of Life Habits  |
| Awareness Questionnaire  |
| Mayo-Portland Adaptability Inventory (MPAI)  |
| Medicare G-Codes   |
| The NIH Toolbox for Assessment of Neurological and Behavioral Function   |
| Neurology quality of life (Neuro-QOL)  |
| Patient-Reported Outcomes Measurement Information System (PROMIS)  |
| Rancho Scales  |